

# **Ticosonde CFH at Costa Rica: A seasonal climatology of tropical UT/LS water vapor and inter-comparisons with MLS and CALIPSO**

**Henry B. Selkirk, GESTAR/NASA GSFC**

with

Holger Vömel, Deutscher Wetterdienst

Melody Avery, NASA LaRC

Karen Rosenlof, NOAA ESRL

Sean Davis and Dale Hurst, CIRES/NOAA ESRL

Mark Schoeberl, STC

Jorge Andrés Díaz, Univ. Costa Rica

Gary Morris, St. Edwards University





Aura STM 2014

# Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder**
- B. Intercomparison to MLS v3.3 WV**
- C. Relationship between sonde saturation and CALIOP cloud fraction**
- D. Changes of WV in the tropics UT/LS**
  - Comparisons of sondes to MLS at 82 hPa
  - 2-km WV time series at Costa Rica





Aura STM 2014

# Science drivers

- A. Processes controlling stratospheric water vapor: local freeze-drying, transport from remote tropical regions, in-mixing from higher latitudes**
- B. Continuing differences among *in situ* water vapor instruments**
- C. *Jensen et al.* [2013] found differing saturation regimes for low- and hi-particle density cirrus**
- D. Radiative forcing of UT/LS water vapor and impact in warming climate**





Aura STM 2014

# Talk topics


- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder**
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS
  - Comparisons of sondes to MLS at 82 hPa
  - 2-km WV time series at Costa Rica






## 2005-2013

## Frostpoint temperature, ambient temperature, and ozone



- ➡ Ozone gradient change nearly coincident with mean coldpoint
- ➡ Prominent inflection in  $O_3$  profile @ 20 km




- ➡ Ozone gradient change well below mean coldpoint
- ➡ Smooth ozone profile throughout





## Second half of year – JJA and SON




- 4 ppmv “hygropause” clearly visible at 468 K in JJA
- 5.5 ppmv maximum rises out of TTL in SON to 415 K



Aura STM 2014

# Tape recorder - II

First half of year – DJF and MAM



- Profile minimum near trop in DJF and slow movement upward through MAM
- But SON max near 420 K has moved up to 460 K by DJF and 525 K in MAM
- Weak ascent in LMS during DJF and MAM, but more rapid above 450 K





Aura STM 2014

# Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV**
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS
  - Comparisons of sondes to MLS at 82 hPa
  - 2-km WV time series at Costa Rica






Aura STM 2014

## Coincidences with MLS v3.3


$\pm 3$  hours,  $\Delta \leq 600$  km



### December – February

65 coincident soundings:

Costa Rica (35), Biak (19), Kototabang (4), Tarawa (4)



### June – September

28 coincident soundings:

Costa Rica (22), San Cristobál (6)





Aura STM 2014

# Talk topics




- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction**
- D. Changes of WV in the tropics UT/LS
  - Comparisons of sondes to MLS at 82 hPa
  - 2-km WV time series at Costa Rica





Aura STM 2014

## $RH_{ice}$ : DJF vs. JJA









Aura STM 2014

# CALIOP Vertical Feature Mask comparison 15 July 2005



VFM cloud analyzed at San José 10.5 to 14.5 km



CFH saturation layer 11.4 to 15 km



- 28 VFM/CFH coincidences closer than 400 km, 2006-2011
- Preliminary finding is that CALIOP is capturing most of the layers > 1 km deep
- However, VFM is just first step – ultimate goal is to compare to ice water content








Aura STM 2014

# CALIOP Mean Cloud Fraction at San Jose, Costa Rica



*Peak frequencies of sonde saturation*







Aura STM 2014


# Talk topics

- A. Seasonal structure at Costa Rica: T, ozone and the water vapor tape recorder
- B. Intercomparison to MLS v3.3 WV
- C. Relationship between sonde saturation and CALIOP cloud fraction
- D. Changes of WV in the tropics UT/LS**
  - Comparisons of sondes to MLS at 82 hPa
  - 2-km WV time series at Costa Rica





## Lower stratospheric water vapor MLS, frostpoint sondes and tropical CPT



- Balloon sonde
- MLS zonal mean, 82 hPA
- MERRA tropical CPT

Adapted from Fig. 2.43, BAMS 2014: State of the Climate 2013






Aura STM 2014

# Time series calculations

- **Costa Rica CFH observations, July 2005 – July 2013**
- **Monthly time series of 2-km thick layers**
  - Linear average of 40 values at 50-m grid points
  - Centered at 3, 5, 7..... km
  - Layer average points computed only if 30 or more grid levels present
- **Multiple soundings in a month composited to a single value**
- **Monthly time means at each 1-km level derived for 8-year period**
  - Minimum sample size of 4 for each monthly level time mean
  - Exclude levels with 6 or more monthly time means missing
  - However, did allow interpolation of up to 2 consecutive missing months
  - Highest level meeting this criterion was 21 km
- **De-seasonalized anomaly time series generated**
  - Subtract long-term means for the month from each value at all levels meeting above criteria
- **Did linear fits in IGOR Pro to anomaly time series to check for trends**










Aura STM 2014

## Layer-average WV time series and de-seasonalized anomalies







Aura STM 2014

# Summary

- Tape recorder structure at Costa Rica consistent with zonal mean, including slow uplift in early part of year in lowermost stratosphere
- Intercomparing MLS and CFH:
  - No real change from *Vömel et al.* [2007]: MLS drier than CFH in UT, wetter in LMS
  - Caveats: UT result swamped by large scatter; difference in LMS within instrumental uncertainties
- Seasonal mean CALIOP cloud fractions consistent with pattern of saturated sondes
  - Preliminary comparisons with coincident CALIOP feature mask data indicate that it can see most saturated sonde layers deeper than about 1 km are
  - However, addressing low- vs. high-density cirrus question will require information on ice water content
- Changes of WV in the tropics UT/LS:
  - Ticosonde CFH following tropical CPT values
  - Weak upward trends over nine years of program not significant – which is not unexpected given dominance of interannual variability